

## Biochemical Constituents and Antioxidant Capacity of Some Seaweeds from Red and Mediterranean Coasts of Egypt

\*M. M. Emam , H. A. Mansour, A. M. Shaaban and N. H. Mostafa

Department of Botany, Faculty of Sciences, Ain Shams university, 11566, Cairo, Egypt.

**P**HYTOCHEMICAL screening was performed on six algal genera belonging to three marine macroalgal divisions (Chlorophyta, Phaeophyta and Rhodophyta) collected from the coastal region of Red and Mediterranean coasts, Egypt during spring (2012). The antioxidant potential of the candidate seaweeds was evaluated by measuring the ferric reducing power (FRAP), total antioxidant capacity (TAC), total phenol and flavonoids, ascorbic acid as well as glutathione contents were determined. Significant individual differences of the biochemical parameters were recorded in all tested marine algae. The macroalgal phytochemical analysis showed quantitative variations in the total soluble and insoluble carbohydrates and proteins as well as the glycerol contents. The highest accumulation of carbohydrate was estimated in the members of Chlorophyta followed by Phaeophyta and Rhodophyta divisions. The protein content was higher in the species of Rhodophyta, moderate in Chlorophyta and the lowest was estimated in the members of Phaeophyta. In addition, the maximum glycerol content was recorded in *Dictyota*. Green algae exhibited greater levels of chlorophylls *a* and *b*. However, among all the studied three groups, the highest carotenoids were estimated in *Padina* followed by *Dictyota*; both belonging to Phaeophyta while the lowest carotenoid content was recorded in *Gracelaria* (Rhodophyta). The highest carbohydrate content was recorded in *Enteromorpha* where the lowest phenol was estimated. Results also revealed that, there was no relation between antioxidant activity and total phenols or flavonoid content. All the studied species are considered to be a rich source of antioxidants (ASA and GSH). Accordingly, these seaweed species have a high anti-oxidative potential and can provide dietary alternatives.

**Keywords:** Antioxidant, Biochemical composition, Ferric reducing power; Seaweeds.

**Abbreviations:** ASA Ascorbic acid, FRAP Ferric reducing ability power, GSH Reduced Glutathione, H<sub>2</sub>O<sub>2</sub> Hydrogen peroxide, ROS reactive oxygen species and TAC Total antioxidant capacity.

---

\*Corresponding author: Manal Emam, Email: [Emammanal@ymail.com](mailto:Emammanal@ymail.com)

Increasing demand for natural antioxidants having less or no side effects is one of the main issues for marine biologists. Edible seaweeds have historically been consumed by coastal populations as a source of pharmacological and food additives with high antioxidant capacity (Rimber and Indy 2007; Gupta and Abu-Ghannam, 2011; Peng *et al.*, 2013 and Rameshkumar *et al.*, 2013). In fact, seaweeds live in complex habitats under extreme conditions (salinity, temperature, nutrients, UV etc.), therefore, the adaptation process produce a great variety of biologically active metabolites, which cannot be found in other organisms (Carlucci, *et al.*, 1999). Seaweeds had been used for many years directly for human consumption, animal feed and as manure because of their high calcium and protein content (Kumari *et al.*, 2010). It is also an ingredient for cosmetics industries and is used as fertilizer (Rupérez, 2002). Interestingly, seaweeds are used in several industries as the gelling, thickening, emulsifying, binding, stabilizing, clarifying and agars (Chopin, 2007). Moreover, macroalgal polysaccharides are used in the food, cosmetics, paints, textile, paper, rubber and building industries. In addition, they are used in medicine and in pharmacology for their antimicrobial, antiviral, antitumor and anticoagulant properties (Fleurence, 1999 and Sabina *et al.*, 2005). Recent studies show the antioxidant abilities of many types of seaweeds (Lim *et al.*, 2002; Park *et al.*, 2005; Duan *et al.*, 2006 and Kuda and Ikemori, 2009). Marine algae, like other photosynthesizing plants, are exposed to different environmental conditions that lead to the generation of reactive oxygen species (Asada, 1999). The injurious effects occurs when the capacity of antioxidant defense system are lower than the generation of reactive oxygen species (ROS). However, the stability of their structural components suggests the presence of protective defense systems (Fujimoto 1990; Matsukawa *et al.*, 1997 and Lim *et al.*, 2002) to alleviate the injurious effects of ROS. Ascorbic acid (ASA) and reduced glutathione (GSH) are responsible for scavenging such ROS by the Hallwell-Asada pathway (Horemans *et al.*, 2000). A major function of ascorbic acid is to scavenge H<sub>2</sub>O<sub>2</sub> through the ascorbate / glutathione pathway (Zhang and Kirkham, 1996). Ascorbate works in co-operation with GSH providing synergetic protection of the membranes.

Natural pigments also have received particular attention as they exhibit various beneficial biological activities such as antioxidant, anticancer, anti-inflammatory, anti-obesity, anti-angiogenic and neuroprotective activities. Therefore, various natural pigments isolated from marine algae have attracted much attention in the fields of food, cosmetic and pharmacology (Pangestuti and Kim, 2011). Estimation of total antioxidant capacity is therefore a valuable tool in evaluation of seaweeds nutritional value (Stajner *et al.*, 2011). In addition to the well known antioxidant system, the light was thrown on the essential role of flavonoids, and phenolic compounds as effective antioxidants (Michalak, 2006). The antioxidant activity of these compounds are mainly attributed to scavenging activity against superoxide and hydroxyl radicals, chelating ability, quenching singlet and triplet oxygen and reducing power (Ruberto, *et al.*, 2001 and Athukorala *et al.*, 2006). In this respect, algae can be considered as an important

source of antioxidant compounds that could be suitable also for protecting our bodies against the reactive oxygen species. Currently, the objective of this study was to evaluate the proximate composition of Egyptian seaweeds (*Sargassum dentifolium*, *Padina pavonia*, *Dictyota dichotoma*, *Gelidium latifolium*, *Gracilaria dura*, *Enteromorpha intestinales* and *Codium tomentosum*) as new antioxidant sources for human and animal nutrition.

### Material and Methods

#### Algal Sampling

During spring 2012, the studied seven seaweeds (marine macro-algae) were collected from Red and Mediterranean coasts of Egypt (Table 1). These including, *Sargassum dentifolium*, *Padina pavonia* and *Dictyota dichotoma* (brown algae belonging to Phaeophyta division) were collected from the coastal region of Red sea (from Hurghida), *Gracilaria dura* (red alga belong to Rhodophyta division) was collected from Faiyd region, whereas *Enteromorpha intestinales* and *Codium tomentosum* (green algae belonging to Chlorophyta division) as well as the red one *Gelidium latifolium* were collected from northern coast of Alexandria (Al-Montazh region). The harvested seaweed samples were immediately washed with seawater to remove the sand particles and epiphytes. Then they were kept in an ice box and immediately transported to the laboratory. After that, the samples were washed thoroughly with tap water to remove the salt and then the species were identified according to Zinova (1967) and Aleem (1993). The samples were spread on blotting paper to remove excess water, then they were divided into two sets, the first lot was air dried and powdered, however the second one was refrigerated at -20 °C until analysis.

TABLE 1. Seaweeds genera collected from coastal region of Egypt.

No.	Seaweeds	Divisions	Sampling site
1	<i>Dictyota dichotoma</i> (hudson) Lamouroux.	Phaeophyta (Brown algae)	Hurghida
2	<i>Padina pavonia</i> (L.) Gaill.		
3	<i>Sargassum dentifolium</i> (Turner) C. Agardh		
4	<i>Gelidium latifolium</i> (Grev.) Born. et. Thur.	Rhodophyta (Red algae)	Al-Montazh
5	<i>Gracilaria dura</i> (C. Ag.) J. Agardh.		Faiyd
6	<i>Enteromorpha intestinales</i> (L.) Link	Chlorophyta (Green algae)	Al-Montazh

#### Chemical analyses

Photosynthetic pigments (Chlorophyll *a*, Chlorophyll *b* and Carotenoids) were extracted in 80% acetone and determined spectrophotometrically as recommended by Metzner *et al.* (1965). Soluble sugars and sugar-free residues were extracted following the method adopted by Homme *et al.* (1992) and

Naguib (1963) respectively. Soluble sugars and those obtained after polysaccharides hydrolysis were estimated using the anthrone reagent as described by Blakeney and Mutton (1980). Moreover, glycerol content was estimated according to Lambart and Neish (1950). Soluble proteins were extracted according to the method described by Guy *et al.* (1992). Meanwhile, the water insoluble residue remaining after extraction of soluble proteins was extracted with 1 N NaOH. Soluble proteins and those resulting after insoluble residue hydrolysis were measured using BIO-RAD protein assay dye reagent according to the method adopted by Bradford (1976).

Antioxidants as glutathione and ascorbic acid were extracted and estimated following the methods described by Smith (1985) and Mukherjee and Choudhuri (1983), respectively. The amount of total phenol was determined with the Folin-Ciocalteu reagent using the method of Malik and Singh (1980) and the absorbance was measured at 765 nm using spectrophotometer (Spectronic 601, Milton Roy Company). Total flavonoid content can be determined by Aluminum chloride method (Harborn, 1998) and the absorbance was measured at 415 nm. Ferric reducing power (FRAP) was estimated according to Dorman *et al.* (2003). Moreover, total antioxidant capacity was determined according to the method of Oyaizu (1986).

#### *Statistic Analysis*

Data were subjected to an analysis of variance, and the means were compared using the least significant difference (LSD) test at the 0.05 level according to the method of Snedecor and Cochran (1982).

### **Results and Discussion**

Totally seven seaweeds were tested for their nutritive properties viz. carbohydrates, proteins, glycerol, flavonoids and total phenols. Measurable differences in nutritional composition were apparent among the candidate macroalgal species. Carbohydrates and proteins are the most important biochemical components in algal biomass. Data presented in (Table 2) showed that, the concentration of carbohydrate was higher in the specie of Chlorophyta followed by Phaeophyta and Rhodophyta respectively. Similarly, Haroon (2000) and Anantharaman *et al.* (2013) recorded highest carbohydrates content in green seaweeds. In Chlorophyta, the highest carbohydrate content ( $p > 0.05$ ) was observed in *Enteromorpha* (192.2 mg g<sup>-1</sup>FW). In Phaeophyta, the highest carbohydrate content ( $p > 0.05$ ) was observed in *Padina* (93.6 mg g<sup>-1</sup>FW). Meanwhile, there was no any significant difference in total carbohydrate content ( $p < 0.05$ ) between *Sargassum* and *Dictyota*. Moreover, the red seaweed *Gracilaria*, recorded the lowest carbohydrate ( $p > 0.05$ ) concentration (20.4 mg g<sup>-1</sup>FW). Such variations may be attributed to species difference and to differences in their metabolic preferences (Pádua *et al.*, 2004).

TABLE 2. Metabolic constituents of some Egyptian seaweeds.

Seaweeds	Total soluble sugars mg g <sup>-1</sup> FW	Total Insoluble Sugars mg g <sup>-1</sup> FW	Total Sugars mg g <sup>-1</sup> FW	Sucrose mg g <sup>-1</sup> FW	Total Soluble Proteins mg g <sup>-1</sup> FW	Total Insoluble Proteins mg g <sup>-1</sup> FW	Total Proteins mg g <sup>-1</sup> FW	Glycerol µmole/g F.W
<b>Phaeophyta</b>								
<i>S. dentifolium</i>	1.54 <sup>e</sup> ±0.01	49.79 <sup>f</sup> ±0.07	51.31 <sup>f</sup> ±0.09	0.87 <sup>d</sup> ±0.2	4.46 <sup>c</sup> ±0.02	4.36 <sup>c</sup> ±0.003	8.82 <sup>d</sup> ±0.02	144.04 <sup>b</sup> ±1.2
<i>P. pavonia</i>	0.87 <sup>g</sup> ±0.03	92.76 <sup>e</sup> ±0.16	93.59 <sup>e</sup> ±0.16	2.82 <sup>bc</sup> ±0.1	1.97 <sup>h</sup> ±0.01	1.86 <sup>c</sup> ±0.017	3.83 <sup>g</sup> ±0.02	148.47 <sup>b</sup> ±2.2
<i>D. dichotoma</i>	1.45 <sup>e,f</sup> ±0.03	49.72 <sup>f</sup> ±0.94	51.21 <sup>f</sup> ±0.96	0.52 <sup>d</sup> ±0.1	2.97 <sup>fg</sup> ±0.08	1.20 <sup>f</sup> ±0.096	4.18 <sup>g</sup> ±0.07	238.51 <sup>a</sup> ±2.9
<b>Rhodophyta</b>								
<i>G. latifolium</i>	5.63 <sup>a</sup> ±0.2	20.85 <sup>g</sup> ±1.4	26.63 <sup>g</sup> ±1.2	2.49 <sup>bc</sup> ±0.3	5.55 <sup>b</sup> ±0.18	4.66 <sup>c</sup> ±0.02	10.21 <sup>b</sup> ±0.16	136.35 <sup>c</sup> ±3
<i>G. dura</i>	3.75 <sup>b</sup> ±0.03	16.72 <sup>g</sup> ±0.67	20.4 <sup>g</sup> ±0.67	3.41 <sup>b</sup> ±0.07	6.42 <sup>a</sup> ±0.01	5.27 <sup>b</sup> ±0.23	11.69 <sup>a</sup> ±0.23	129.49 <sup>d</sup> ±0.9
<b>Chlorophyta</b>								
<i>E.intestinales</i>	1.78 <sup>d</sup> ±0.01	190.5 <sup>c</sup> ±6.08	192.2 <sup>c</sup> ±6.08	3.48 <sup>b</sup> ±0.1	3.40 <sup>c</sup> ±0.15	6.13 <sup>a</sup> ±0.11	9.53 <sup>c</sup> ±0.21	52.23 <sup>f</sup> ±0.4
LSD (>0.05)	species	0.234	17.219	17.161	1.132	0.216	0.33	5.962
D (0.05)	Division	0.128	9.431	9.399	0.62	0.118	0.181	3.265

All the values are mean ± SE. Column wise values with same superscripts of this type indicate no significant difference (P > 0.05).

The potential production of glycerol by seaweeds is of obvious economic benefit (Ben-Amotz and Avron, 1980; Chen and Chi, 1981). Most seaweeds produce glycerol exclusively by degradation of starch. This is the case in our study; the glycerol content was in inverse to the insoluble carbohydrate content. The maximum (p>0.05) glycerol content was recorded in the members of phaeophyta where the lowest one was measured in Chlorophyta specie (Table 2). It is interesting to note that, there was no significant difference in glycerol content (p<0.05) between *S. dentifolium* and *P. pavonia*. In seaweeds, carbohydrate metabolism may be shifted toward glycerol production (Avron, 1992). Similarly, Bentalel *al.*, (1990) on his study on *Dunaliella*, suggested a dynamic inter-conversion between the two major carbon pools (glycerol and starch) to meet the osmotic requirements.

Like carbohydrates and glycerol, there was a significant differences ( $p > 0.05$ ) of protein contents among all studied seaweeds. In the present study, protein content showed remarkable variation with highest values in Rhodophyta followed by Chlorophyta and Phaeophyta. Protein content varied from 3.8 to 11.6 mg g<sup>-1</sup>FW; maximum protein ( $p > 0.05$ ) was recorded in *Gracilaria* (11.6 mg g<sup>-1</sup>FW) followed by *Geledium* (10.2 mg g<sup>-1</sup>FW) and *Enteromorpha* (9.5 mg g<sup>-1</sup>FW). The minimum protein ( $p > 0.05$ ) concentration was estimated in *Padina* (3.8 mg g<sup>-1</sup>FW) followed by *Dictyota* (4.1 mg g<sup>-1</sup>FW) from Phaeophyta (Table 2). Nevertheless, higher protein contents were recorded in red and green algae; the protein content of brown algae is generally low. The findings of the present study coincided well with Dere *et al.*, (2003) and Rameshkumar *et al.* (2013) who reported a similar trend of higher protein content in species of Rhodophyta, moderate in Chlorophyta and the lowest in Phaeophyta. According to the study of Marinho-Soriano *et al.* (2006), the high carbohydrate content was concomitant with decreasing the nitrogen and proteins content. This is the case in our study; the higher protein content in red algae which recorded the lowest carbohydrates. Protein content varied among different genera and also in different species of the same genus (Dhargalkar *et al.*, 1980). On the contrary, Burtin (2003) estimated high protein content in green seaweeds. However, Wong and Cheung (2001) demonstrated that protein content of the red seaweeds was significantly higher than those of green seaweeds.

The environment in which seaweeds grow is harsh and can lead to the formation of strong oxidizing agents but seaweeds seldom suffer any serious photodynamic damage during metabolism. Such fact implies that seaweed cells have evolved complex protective system (Matsukawa *et al.*, 1997) to tolerate salinity. Attention of researchers was focused on the ability of algae to accumulate antioxidant compounds. Heo *et al.* (2005) identified the potential antioxidant compounds as some pigments (e.g. chlorophyll, carotenoid) and polyphenols (e.g. phenolic acid, flavonoid). Data presented in Table 3 indicated that, amongst all the species of the three algal groups, the order of chlorophyll *b* content is as follows: Chlorophyta > Phaeophyta > Rhodophyta. The results of the current study corroborated with the findings of Francisco *et al.* (2006) and Kumar *et al.* (2009) stating highest chlorophyll *a* and *b* contents in green algae and lowest in red algae. Among all the three groups, highest carotenoids ( $p > 0.05$ ) were reported in *Padina* (6.62 ug g<sup>-1</sup> FW) followed by *Dictyota* (5.56 ug g<sup>-1</sup> FW), both belonging to the Phaeophyta while the lowest carotenoid content ( $p > 0.05$ ) was observed in *Gracelaria* (1.3 ug g<sup>-1</sup> FW) belonging to the Rhodophyta. Similar results were reported by Fritsch (1971) and Dere *et al.* (2003). Indeed, chlorophyll *a* exhibited antioxidant activity (Endo *et al.*, 1985), act as potent synergist of vitamin E (Mendiola *et al.*, 2005). Carotenoids are known to quench the excited molecules and singlet oxygen (Bondarev, 1997). In this study, the higher antioxidant activity of *Dictyota* (Table 3) might be attributed to the high content of the lipid soluble total chlorophylls especially Chl *a* (Stajner *et al.*, 1999) and / or to the high content of total carotenoids (Rivero *et al.*, 2003).

TABLE 3. Chlorophyll *a* and *b*, carotenoid contents of some Egyptian seaweeds.

Seaweeds	Chlorophyll a	Chlorophyll b	Carotenoids	Ch <sub>a</sub> /Ch <sub>b</sub>	Total Pigments
	µg/g FW				
<b>Phaeophyta</b>					
<i>S. dentifolium</i>	7.50 <sup>fg</sup> ±0.5	2.64 <sup>fg</sup> ±0.06	4.14 <sup>c</sup> ±0.3	2.87 <sup>b</sup> ±0.3	14.28 <sup>c</sup> ±1
<i>P. pavonia</i>	10.07 <sup>c</sup> ±0.6	3.55 <sup>de</sup> ±0.1	6.62 <sup>a</sup> ±0.5	2.85 <sup>b</sup> ±0.05	20.24 <sup>b</sup> ±0.8
<i>D. dichotoma</i>	10.17 <sup>de</sup> ±0.2	2.89 <sup>ef</sup> ±0.01	5.56 <sup>b</sup> ±0.07	3.52 <sup>a</sup> ±0.1	18.62 <sup>b</sup> ±0.1
<b>Rhodophyta</b>					
<i>G. latifolium</i>	2.94 <sup>f</sup> ±0.3	1.94 <sup>h</sup> ±0.01	1.4 <sup>fg</sup> ±0.08	1.52 <sup>c</sup> ±0.09	6.28 <sup>f</sup> ±0.2
<i>G. dura</i>	0.1±2.36	2.07 <sup>gh</sup> ±0.1	1.32 <sup>g</sup> ±0.03	1.16 <sup>cd</sup> ±0.2	5.75 <sup>f</sup> ±0.2
<b>Chlorophyta</b>					
<i>E. intestinales</i>	10.65 <sup>de</sup> ±1.1	9.44 <sup>a</sup> ±0.5	3.02 <sup>d</sup> ±0.05	1.13 <sup>cd</sup> ±0.01	23.12 <sup>a</sup> ±1.3
<b>LSD</b>	0.560	0.673	0.622	0.413	2.069
<b>(&gt;0.05)</b>					
<b>Species</b>					
<b>Division</b>	00.23	0.369	0.341	0.226	1.134

All the values are mean ± SE. Column wise values with same superscripts of this type indicate no significant difference (P > 0.05).

Phenols are the aromatic metabolites which trigger various biochemical processes of the organisms. Phenolic compounds are known to act as antioxidants not only because of their ability to donate hydrogen atoms or electrons but also because of their stable radical intermediates, which prevent the oxidation of various ingredients, particularly fatty acids and oils (Yuan *et al.*, 2005). Data presented in Table 4 revealed that, the total phenol was in inverse to carbohydrate content in most species under investigation. The highest carbohydrate content (p>0.05) was recorded in *Enteromorpha* (19c mg g<sup>-1</sup>FW) where the lowest phenol (0.34 mg g<sup>-1</sup>FW) was estimated. The carbohydrate metabolism might be channeled for the production of total phenols (Tongnetti and Johnson, 1999), thus explaining the reason why the production of secondary metabolites was up-regulated with decreasing carbohydrates. Moreover, data in Table 4 indicated that, there was no significant difference in total phenols content (p<0.05) between green algae members. Likewise, the highest flavonoid contents were showed in the investigated macro-algal species of Phaeophyta followed by Chlorophyta and Rhodophyta respectively. In the members of the Phaeophyta the highest flavonoid contents (p>0.05) were 5.4 mg g<sup>-1</sup> DW, 3.4 mg g<sup>-1</sup> D.W. and 3.2 mg g<sup>-1</sup> DW in *Padina*, *Dictyota* and *Sargassum* respectively. The antioxidative properties of flavonoids are due to several different mechanisms, such as scavenging of free radicals, chelating of metal ions, such as iron and copper, and inhibition of enzymes responsible for free radical generation. Depending on their structure, flavonoids are able to scavenge practically all known ROS (Chanda and Dave, 2009). Previous study gave similar results and explained that, there was no correlation between antioxidant activity and total phenol or flavonoid content (Yu *et al.* 2003; Dasgupta, 2007; Alam *et al.*, 2012).

Data presented in Table 4 also indicated variations in TAC which may be attributed to species differences. Accordingly, all the studied species are considered to be a rich source of antioxidants. The antioxidant activity of the plant extract has been found to be associated with the reducing power property. Among the seven species studied, *Sargassum* had the highest concentration of ascorbic acid whereas *padina* had the lowest. *Sargassum* also had the richest source of glutathione. Furthermore, the highest reducing power ( $p>0.05$ ) was observed in *Dictyota* ( $1.04 \mu\text{g g}^{-1}$  DW) and the lowest reducing power was estimated in the green algae *Enteromorpha* ( $0.19 \mu\text{g g}^{-1}$  DW). Interestingly, *Dictyota* also exhibited the highest antioxidant activity suggesting the relationship between antioxidant activities and reducing power (Table 4). Seenivasan (2013) reported that brown seaweeds exhibited good antioxidant activity when compared to red and green seaweeds. Indeed, all species under studies showed high reducing power and the studied seaweed extracts were able to reduce  $\text{Fe}^{+3}$  to  $\text{Fe}^{+2}$  as a function of reducing power.

In conclusion, the present study evident that, the seaweeds under investigation (*Sargassum dentifolium*, *Padina pavonia*, *Dictyota dichotoma*, *Gelidium latifolium*, *Gracilaria dura* and *Enteromorpha intestinales*) are interesting natural source of some antioxidant compounds that could be used as functional ingredients and can provide dietary alternatives. Moreover, further research is underway for isolation and characterization of natural antioxidant having less or no side effects, for use in foods or medicines to replace synthetic antioxidant.

**TABLE 4. Antioxidant capacity, ferric reducing power, ascorbic acid, glutathione, total phenols and flavonoid contents of some Egyptian seaweeds.**

Seaweeds	Ascorbic Acid $\mu\text{Mole/g}$ F.W	Glutathione $\text{mMole/g}$ F.W	Total Phenols $\text{mg/g}$ F.W	Total Flavonoids $\text{mg/g}$ D.W	Reducing Power $\mu\text{g/g}$ D.W	Total Antioxidant Capacity $\text{mg/g}$ D.W
<b>Phaeophyta</b>						
<i>S. dentifolium</i>	0.70 <sup>a</sup> ±0	13.17 <sup>a</sup> ±0.6	0.82 <sup>b</sup> ±0.06	3.21 <sup>de</sup> ±0.09	0.20 <sup>e</sup> ±0.01	189.08 <sup>bc</sup> ±1.3
<i>P. pavonia</i>	0.27 <sup>d</sup> ±12.17	11.13 <sup>b</sup> ±1	0.77 <sup>b</sup> ±0.04	5.43 <sup>b</sup> ±0.05	0.46 <sup>b</sup> ±0.02	165.51 <sup>bc</sup> ±3.6
<i>D. dichotoma</i>	0.47 <sup>b</sup> ±15.68	8.47 <sup>cd</sup> ±0.2	0.61 <sup>bc</sup> ±0.09	3.42 <sup>d</sup> ±0.1	1.04 <sup>a</sup> ±0.05	1471.26 <sup>a</sup> ±123.5
<b>Rhodophyta</b>						
<i>G. latifolium</i>	0.31 <sup>cd</sup> ±1.42	8.22 <sup>de</sup> ±0.5	1.21 <sup>a</sup> ±0.02	2.37 <sup>f</sup> ±0.1	0.31 <sup>cd</sup> ±0.02	249.42 <sup>b</sup> ±9.9
<i>G. dura</i>	0.33 <sup>c</sup> ±13.58	8.42 <sup>cd</sup> ±0.3	1.08 <sup>a</sup> ±0.07	1.79 <sup>e</sup> ±0.1	0.22 <sup>de</sup> ±0.01	194.82 <sup>bc</sup> ±11.4
<b>Chlorophyta</b>						
<i>E.intestinales</i>	0.45 <sup>b</sup> ±37.9	7.73 <sup>e</sup> ±0.2	0.39 <sup>cd</sup> ±0.03	3.83 <sup>c</sup> ±0.2	0.19 <sup>e</sup> ±0.01	134.86 <sup>bc</sup> ±8.3
<b>LSD (0.05&lt;)</b>						
Species	0.045	1.479	2.258	0.359	0.093	118.67
Division	0.025	0.81	1.236	0.196	0.051	64.99

All the values are mean ± SE. Column wise values with same superscripts of this type indicate no significant difference ( $P > 0.05$ ).



## References

- Alam, M. N., Wahed, T. B., Sultan, F., Ahmed, J., and Hasan, M. (2012)** *In vitro* antioxidant potential of the methanolic extract of *Bacopa monnieri* L., *Turk. J. Pharm. Sci.*, **3**, 285-292.
- Aleem, A.A. (1993)** The Marine Algae of Alexandria, Egypt.
- Anantharaman, P., Parthiban, C., Saranya, C., Girija, K., Hemalatha, A. and Suresh, M., (2013)** Biochemical composition of some selected seaweeds from Tuticorin coast. *Adv. Appl. Sci. Res.*, **4**, 362-366.
- Asada, K. (1999)** The water-water cycle in chloroplasts: Scavenging of active oxygen and dissipation of excess photons. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, **50**, 601-639.
- Athukorala, Y., Nam, K. and Jeon, Y. (2006)** Antiproliferative and antioxidant properties of an enzymatic hydrolysate from brown alga *Ecklonia cava*. *Food Chem. Toxicol.*, **44**, 1065-1074.
- Avron, M. (1992)** Osmoregulation. In: *Dunaliella: "Physiology, Biochemistry, and Biotechnology"*, M Avron and A Ben-Amotz (Ed.), CRC Press, Inc., Boca Raton, Florida, pp.135-164.
- Ben-Amotz, A. and Avron, M. (1980)** In: *"Glycerol, J3-Carotene and Dry Algal Meal Production by Commercial Cultivation of Dunaliella"*. Algae Biomass, G Shelef and CJ Soeder (Ed.), Elsevier/North-Holland Biomedical Press, Amsterdam.
- Bental, M., Pick, U., Avron, M. and Degani, H. (1990)** The Role of Intracellular Orthophosphate in Triggering Osmoregulation in the Alga *Dunaliella salina*. *Europ. J. of Biochem.*, **188**, 117-122.
- Blakeney, A.B. and Mutton, L.L. (1980)** A simple colorimetric method for the determination of sugars in fruits and vegetables. *J. Sci. Food Agri.*, **31**, 889-897.
- Bondarev, S.L. (1997)** Photophysics of  $\beta$ -carotene and related compounds. *J. Appl. Spectro.*, **64**, 1-5.
- Bradford, M. M. (1976)** A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein dye binding. *Anal. Biochem.*, **72**, 248-254.
- Burtin, P. (2003)** Nutritional value of seaweeds. *Elect. J. Environ. Agric. Food Chemist.*, **2**, 498-503.
- Carlucci, M. J., Scolaro, L. A. and Damonte, E. B. (1999)** Inhibitory action of natural carrageenans on Herpes simplex virus infection of mouse astrocytes. *Chemother.*, **45**, 429-36.
- Chanda, S. and Dave, R. (2009)** *In vitro* models for antioxidant activity evaluation and some medicinal plants possessing antioxidants properties: An overview, *Afri. J. of Microbiol. Res.*, **3**, 981-96.

- Chen, B.J and Chi, C.H. (1981)** "Process Development and Evaluation for Algal Glycerol Production. *Biotechnol. and Bio-engineer.*", **23**, 1267-1287.
- Chopin, T. (2007)** Closing remarks of the new president of the international seaweed Association. "The 19<sup>th</sup> International Seaweed Symposium Kobe-Japan".
- Dasgupta, N., De, B., (2007)** Antioxidant activity of some leafy vegetables of India: A comparative study, *Food Chem.*, **101**, 471-474.
- Dere, S., Dalkiran, N., Karacaoglu, D., Gamze Yildiz, G. and Dere, E. (2003)** The determination of total protein, total soluble carbohydrate and pigment contents of some macro algae collected from Gemlik-Karacaali (Bursa) and Erdek-Ormanli (Balikesir) in the Sea of Marmara, *Turkey. Oceanologia*, **45**, 453-471.
- Dhargalkar, V. K., Jagtap, T.G. and Untawale, A.G. (1980)** Biochemical constituents of seaweeds along the Maharashtra coast. *Indian J. Mar. Sci.*, **9**, 297-299.
- Dorman, H., Kosar, M., Kahlos, K., Holm, Y. and Hiltunen, R. (2003)** Antioxidant properties and composition of aqueous extracts from *Mentha* species, hybrids, varieties, and cultivars. *J. of Agri. and Food Chem.*, **51**, 4563-4569.
- Duan, X. J., Zhang, W. W., Li, X. M. and Wang, B. G. (2006)** Evaluation of antioxidant property of extract and fractions obtained from a red alga, *Polysiphonia urceolata*. *Food Chem.*, **95**, 37-43.
- Endo, Y., Usuki, R. and Kaneda, T. (1985)** Antioxidant effects of chlorophyll and pheophytin on the autooxidation of oils in the dark. II. The mechanism of antioxidative action of chlorophyll. *J. American Oil Chem. Soc.*, **62**, 1387-90.
- Fleurence, J. (1999)** Seaweed proteins: Biochemical, nutritional aspects and potential uses. *Trends Food Sci. Tech.*, **10**, 25-28.
- Francisco, J. L. G., Aguilera, J. and Jimenez, C. (2006)** The response of nutrient assimilation and biochemical composition of Arctic sea-weeds to a nutrient input in summer. *J. Exp. Bot.*, **57**, 2661-2671.
- Fritsch, F. E. (1971)** "The Structure and Reproduction of the Algae", Vol I, Cambridge Univ. Press, Cambridge. Pp 791 .
- Fujimoto, K. (1990)** Antioxidant activity of algal extracts. In: " *Introduction to Applied Phycology*". Akatsuka I (Ed.), SPB Academic Publishing, The Hague. pp 199-208.
- Gupta, S. and Abu-Ghannam, N. (2011)** Bioactive potential and possible health effects of edible brown seaweeds. *Trends Food Sci. Technol.*, **22**, 315-326.
- Guy, C.L., D. Haskell, L. Neven, P. Klein and C. Smelser. (1992)** Hydration state-responsive protein like cold and drought stress in spinach. *Planta*, **188**, 265-270.
- Haroon, M. (2000)** The biochemical composition of *Enteromorpha spp.* From the Gulf of Gdansk coast on the southern Baltic sea. *Oceanologia*, **42**, 19-28.

- Harborne, J. B. (1998)** "Photochemical Methods: A Guide to Modern Technique of Plant Analysis", 3<sup>rd</sup> ed. Chapman and Hall, London, UK, p. 302.
- Heo, S. J., Park, E. J., Lee, K.W. and Jeon, Y. J. (2005)** Antioxidant activities of enzymatic extracts from brown seaweeds. *Biores. Technol.*, **96**, 1613–1623.
- Homme, P. M., Gonzalez, B. and Billard, J. (1992)** Carbohydrate content, fructane and sucrose enzyme activities in roots, stubble and leaves of rye grass (*Lolium perenne* L.) as affected by source/sink modification after cutting. *J. Plant Physiol.*, **140**, 282-291.
- Horemans, N., Foyer, C.H., Potters, G. and Asard, H. (2000)** Ascorbate function and associated transport systems in plants. *Plant Physiol. Biochem.*, **38**, 531.
- Kuda, T. and Ikemori, T. (2009)** Minerals, polysaccharides and antioxidant properties of aqueous solutions obtained from macroalgal beach-coasts in the Noto Peninsula, Ishikawa, Japan. *Food Chem.*, **112**, 575-581.
- Kumar, N.J.I., Kumar R.N., Bora A., Kaur Amb, M. and Chakraborty, S. (2009)** An evaluation of pigment composition of eighteen marine macro algae collected from Okha coast, Gulf of Kutch, India. *Our Nature*, **7**, 48-55.
- Kumari, P., Kumar, M., Gupta, V., Reddy, C.R.K. and Jha, B. (2010)** Tropical marine macroalgae as potential sources of nutritionally important PUFAs. *Food Chemist.*, **120**, 749-757.
- Lambert, M. and Neish A.C. (1950)** Rapid method for estimation of glycerol in fermentation solutions. *Can. J. Res.*, **28**, 83-89.
- Lim, S. N., Cheung, P.C.K., Ooi V.E.C. and Ang, P.O. (2002)** Evaluation of antioxidative activity of extracts from a brown seaweed, *Sargassum siliquastrum*. *J. of Agri. and Food Chem.*, **50**, 3862-3866.
- Malik, C. P. and Singh, M. B. (1980)** In: "Plant Enzymology and Histo-Enzymology". Kalyani Publishers. New Delhi. pp. 286.
- Marinho-Soriano, E., Fonseca, P.C., Carneiro, M.A.A., and Moreira, W.S.C. (2006)** Seasonal variation in the chemical composition of two tropical seaweeds. *Bioresour Technol.*, **97**, 2402-2406.
- Matsukawa, R., Dubinsky, Z., Kishimoto, E., Masaki, K., Masuda, Takeuchi, T., Chihara, M., Yamamoto, Y., Niki, E. and Karube, I. (1997)** A comparison of screening methods for antioxidant activity in seaweeds. *J. Appl. Phycol.*, **9**, 29–35.
- Mendiola, J.A., Marin, F.R., Hernandez, S.F., Arredondo, B.O., Senorans, F.J., Ibanez, E. and Reglero, G. (2005)** Characterization via liquid chromatography coupled to diode array detector and tandem mass spectrometry of supercritical fluid antioxidant extracts of *Spirulina platensis* microalga. *J. Separation Sci.*, **28**, 1031–38.
- Metzner, H., Rau, H. and Senger, H. (1965)** Untersuchungen Zur Synchronisierbarkeit einzelner-pigment. Mangol Mutanten Von *Chloella*. *Planta*, **65**, 186-194.

- Michalak, A. (2006)** Phenolic compounds and their antioxidant activity in plants growing under heavy metal stress. *Pol. J. Environ. Stud.*, **15**, 523-530.
- Mukherjee, S. P. and Choudhuri, M. A. (1983)** Implication of water stress-induced changes in the level of endogenous ascorbic acid and hydrogen peroxide in Vigna seedling. *Physiol. Plant.*, **58**, 166-170.
- Naguib, M. I. (1963)** Colourimetric estimation of plant polysaccharides. *Zucher*, **16**, 15-18.
- Oyaizu, M. (1986)** Studies on product of browning reaction prepared from glucose amine. *Jpn. J. Nutr.*, **44**, 307-15.
- Pádua, M. D., Fontoura, P. S. G and Mathias, A. B. (2004)** Chemical composition of *ulvaria oxysperma* (Kützinger) Bliding, *Ulva lactuca* (Linnaeus) and *Ulva fasciata* (Delile) Barazil. *Arch. of Biol. and Technol.*, **47**, 49-55.
- Pangestuti, R and Kim, S. K. (2011)** Biological activities and health benefit effects of natural pigments derived from marine algae. *J. Func. Food.*, **3**, 255-266.
- Park, P. J., Heo, S. J., Park, E.J., Kim, S.K., Byun, H.G., Jeon, B.T. and Jeon, Y.J. (2005)**. Reactive oxygen scavenging effect of enzymatic extracts from *Sargassum thunbergii*. *J. Agric. Food Chem.*, **53**: 6666-6672.
- Peng, Y., Xie, E., Zheng, K., Fredimoses, M., Yang, X., Zhou, X., Wang, Y., Yang, B., Lin, X., Liu, J., and Yonghong Liu, Y., (2013)** Nutritional and chemical composition and antiviral activity of cultivated seaweed *Sargassum naozhouense* Tseng et Lu Mar. *Drugs*, **11**, 20-32.
- Rameshkumar, S., Ramakritinan, C. M. and Yokeshbabu, M. (2013)** proximate composition of some selected seaweeds from Palk bay and Gulf of Mannar, Tamilnadu, India. *Asian J. of Biomed. and Pharma. l Sci.*, **3**, 1-5.
- Rimber, Ir and Indy (2007)** Why is seaweed so important? *M.Sc Thesis* Faculty of Fisheries and Marine Sciences, Sam Ratulangi University, Jln. Kampus Bahu, Manado 95115, Indonesia.
- Rivero, F., Fallarero, A., Castaneda, O., Dajas, F., Manta, E., Areces, F., Mancini, J. Filho and Vidal, A. (2003)** Antioxidant activity *in vivo* and *in vitro* of *Halimeda incrustata* aqueous extracts. *Ciencia-e- Tecnologia-de-Alimentos*, **23**, 256-63.
- Ruberto, G., Baratta, M. T. Biondi, D. M. and Amico, V. (2001)** Antioxidant activity of extracts of the marine algal genus *Cystoseira* in a micellar model system. *J. Appl. Phycol.*, **13**, 403-407.
- Rupérez, P. (2002)** Mineral content of edible marine seaweeds. *Food Chemist.*, **79**, 23-26.
- Sabina, H., Tasneem, S., Kausar, S.Y., Choudhary, M.I. and Aliya, R. (2005)** Antileishmanial activity in crude extract of various seaweeds from the coast of Karachi, Pakistan. *Pakis. J. of Bot.*, **37**, 163-168.

- Seenivasan, R. I. (2013)** *In vitro* antioxidant activity of selected seaweeds from South East coast of India. *Int. J. Pharm, Pharm. Sci.*, **5**, 474-484.
- Snedecor, G. W. and Cochran, W. G. (1982)** "*Statistical Methods*". Ames, Iowa, USA: The Iowa State Univ. Press.
- Smith, I. K. (1985)** Stimulation of glutathione synthesis in photorespiring plants by catalase inhibitors. *Plant Physiol.*, **79**, 1044-1047.
- Štajner, D., Milic N. and Canadanovic-Brunet, J. (1999)** An investigation into the antioxidant activity of *Allium nutans* L. *Phytother. Res.*, **13**, 333-6.
- Štajner, D., Orlović, S., Popović, B. M., Kebert, M. and Galić, Z. (2011)** Screening of drought oxidative stress tolerance in Serbian melliferous plant species *Afr. J. Biotech.*, **10**, 1609-1614.
- Tognetti, R. and Johnson, J. D. (1999)** The effect of elevated atmospheric CO<sub>2</sub> concentration and nutrients supply as gas exchange, carbohydrates and foliar phenolics concentration in live oak (*Quercus virginiana* Mill.) seedlings. *Ann. For. Sci.*, **56**, 379-389.
- Wong, K. H. and Cheung, P. C. K. (2001)** Nutritional evaluation of some subtropical red and green seaweed. Part II. *In vitro* protein digestibility and amino acid profiles of protein concentrate. *Food Chem.*, **72**, 11-17.
- Yu, P. Z., Liu, X. G., Zhou, G. F., Zhang, Q. B. and Li, P.C. (2003)**. Anti hyperlipidemic effects of different molecular weight sulfated polysaccharides from *Ulva pertuse* (chlorophyta). *Pharmacological Res.*, **48**, 543-549.
- Yuan, Y. V, Bone, D. E and Carrington, M. F (2005)** Antioxidant activity of dulse (*Palmaria palmata*) extract evaluated *in vitro*. *Food Chem.*, **91**, 485-494.
- Zhang, J. and Kirkham, M. B. (1996)** Lipid peroxidation in sorghum and sunflower seedlings as affected by ascorbic acid, benzoic acid, and propyl gallate. *J. Plant Phys.*, **149**, 489-493.
- Zinova, A. D. (1967)** Key of green, brown and red algae of southern seas of USSR. Prin. Nauka ANVSSR. Moscow.

(Received 2 / 4/2014;  
accepted 19/5/2014)

## المكونات البيوكيميائية وقدرة مضادات الأكسدة لبعض أنواع الطحالب البحرية من سواحل البحر الأحمر والأبيض المتوسط لمصر.

منال محمد امام، هدي انور منصور، عبد السلام شعبان و نعمات حسن مصطفي  
قسم النبات - كلية العلوم - جامعة عين شمس - القاهرة - مصر.

تهدف هذه الدراسة الى تقييم التركيب الكيميائي والقدرة الكلية لمضادات الأكسدة لستة أنواع من الطحالب البحرية من المنطقة الساحلية في مصر خلال ربيع 2012 ولقد أسفرت النتائج عن وجود اختلافات كمية في تراكم المواد الكربوهيدراتية والبروتينية وكذلك بعض مضادات الأكسدة مثل حمض الاسكوربيك، الجلوتاثيون، الفينولات وكذلك الفلافينويدات.

حيث أظهرت النتائج زيادة معنوية واضحة في تراكم المواد الكربوهيدراتية والبروتينية في الطحالب الخضراء يليها الطحالب البنية ثم الحمراء. بالإضافة الي ذلك تم تسجيل أقصى محتوى للجلسرين في *Dictyota* بينما كان أدنى مستوي له في *Enteromorpha* ومما هو جدير بالذكر أن الطحالب الخضراء أظهرت أعلى مستويات من الكلوروفيل (أ) و (ب) بين المجموعات الثلاث، وسجلت أعلى قيمة للكاروتينيدات في *Padina* تليها *Dictyota*، وكلاهما ينتميان إلى الطحالب البنية بينما سجلت أدنى قيمة في *Gracilaria* (الطحالب الحمراء). ولقد لوحظ وجود علاقة عكسية بين تراكم الفينولات و المواد الكربوهيدراتية في المجموعات الثلاثة. وكشفت النتائج أيضا أنه لم تثبت العلاقة بين القدرة الكلية لمضادات الأكسدة من ناحية و الفينولات و محتوى الفلافونويد من ناحية أخرى. وتبين من النتائج أن جميع الأنواع قيد الدراسة كانت مصدرا غنيا لمضادات الأكسدة مثل حمض الاسكوربيك و الجلوتاثيون (ASA و GSH).

وخلاصة القول، إن هذه الأنواع من الأعشاب البحرية أعطت نتائج واعدة خاصة كمضادات للأكسدة ويمكن أن تستعمل كبداية غذائية.